




1




Collision Physics

Oregon State University
Mike Bailey
mjb@cs.oregonstate.edu



This work is licensed under a [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License](https://creativecommons.org/licenses/by-nc-nd/4.0/).

collision-physics.pptx mjb - July 31, 2021

1


2

There are some cases when you can compute the collision in closed-form

We need to answer the question:

What will cause the ball to stop flying freely through the air?

The answer is – it will hit a wall or the floor. From our rules of physics, the next bounce will happen at one of the solutions to these equations:




$$XRIGHT - RADIUS = x + v_x t$$

$$XLEFT + RADIUS = x + v_x t$$

$$YBOTTOM + RADIUS = y + v_y t + \frac{1}{2} g t^2$$

Note: $g < 0$.



Oregon State University Computer Graphics mjb - July 31, 2021

2

3

There are some cases when you can compute the collision in closed-form

The next bounce will happen at one of the solutions to these equations:

$XRIGHT - RADIUS = x + v_x t \rightarrow t_1 = \frac{XRIGHT - RADIUS - x}{v_x}$

$XLEFT + RADIUS = x + v_x t \rightarrow t_2 = \frac{XLEFT + RADIUS - x}{v_x}$

$YBOTTOM + RADIUS = y + v_y t + \frac{1}{2} g t^2$

$\frac{1}{2} g t^2 + v_y t + (y - YBOTTOM - RADIUS) = 0 \rightarrow t_3, t_4 = \frac{-v_y \pm \sqrt{v_y^2 - 2g(y - YBOTTOM - RADIUS)}}{g}$



$$t_1 = \frac{XRIGHT - RADIUS - x}{v_x}$$

$$t_2 = \frac{XLEFT + RADIUS - x}{v_x}$$

$$t_3, t_4 = \frac{-v_y \pm \sqrt{v_y^2 - 2g(y - YBOTTOM - RADIUS)}}{g}$$

How do you know which of the 4 t values is the one that you should use?

Use the time that is the smallest positive number.

Oregon State University Computer Graphics mjb - July 31, 2021

3

4

There are some cases when you can compute the collision in closed-form

The next bounce will happen at one of the solutions to these equations:

$$t_1 = \frac{XRIGHT - RADIUS - x}{v_x}$$

$$t_2 = \frac{XLEFT + RADIUS - x}{v_x}$$



$$t_3, t_4 = \frac{-v_y \pm \sqrt{v_y^2 - 2g(y - YBOTTOM - RADIUS)}}{g}$$

How do you know which of the 4 t values is the one that you should use?

Use the time that is the smallest positive number.

We want positive, because negative time values happened ago.

We want smallest because we are interested in the collision that happens **first**.

Oregon State University Computer Graphics mjb - July 31, 2021

4

The Physics of Collisions -- Definitions 5

Line of Impact

If the objects' velocities are parallel to the Line of Impact, this is a **Direct Impact**

If the objects' velocities are perpendicular to the Line of Impact, this is a **Tangential Impact**

A combination of the two is called an **Oblique Impact**

Direct Impact

Tangential Impact

Oblique Impact
(both direct and tangential)

Computer Graphics mjb - July 31, 2021

5

The Physics of Collisions – Fundamental Quantities 6

The **momentum** of an object is defined as its mass multiplied by its velocity:

$$\text{Momentum} = mv$$

The **energy** of an object is defined as one half of its mass multiplied by its velocity squared:

$$\text{Energy} = \frac{1}{2}mv^2$$

Oregon State University
Computer Graphics

mjb - July 31, 2021

6

The Physics of Collisions – Conservation of Momentum 7

In a collision, the total momentum after the impact is equal to the total momentum before the impact. Always.

$m_a v_a + m_b v_b = m_a v'_a + m_b v'_b$

where the primes ' refer to velocities after the impact

This is referred to as the **Conservation of Momentum Law**

Momentum is always conserved through **any** collision

Oregon State University
Computer Graphics

mjb - July 31, 2021

7

Conservation of Momentum as Explained by Newton's Cradle 8

http://www.grand-illusions.com/acatalog/Newtons_Cradle.html

<http://dsc.discovery.com/videos/mythbusters-newtons-crane-cradle/>

And, of course, where would any topic be without kittens? ☺

University Computer Graphics

<http://www.buzzfeed.com/koolnewsblog/cats-for-the-love-of-physics-1dgl>

mjb - July 31, 2021


8

The Physics of Collisions – Coefficient of Restitution

9


In a collision, energy is conserved in the entire system, but not necessarily in the form of velocities. (It can become permanent deformation, heat, light, etc.)

This loss of velocity is expressed as the **Coefficient of Restitution** (COR). The COR, e , is how much less the relative velocities of the objects are after impact than they were before impact:



$$v'_b - v'_a = -e(v_b - v_a)$$

(the negative sign is there to indicate the "bounce")




mjb - July 31, 2021

9

The Physics of Collisions – Combining Momentum and Restitution Laws

10



Starting with these two equations:


$$m_a v_a + m_b v_b = m_a v'_a + m_b v'_b$$

$$v'_b - v'_a = -e(v_b - v_a)$$

Treat the two initial velocities as inputs and solve for the two resulting velocities. This gives:

$$v'_a = \frac{m_a v_a + m_b v_b + e m_b (v_b - v_a)}{m_a + m_b}$$

$$v'_b = \frac{m_a v_a + m_b v_b - e m_a (v_b - v_a)}{m_a + m_b}$$



mjb - July 31, 2021

10

The Physics of Collisions with Immoveable Objects

11

To treat the case of mass b being an immoveable object, such as the ground or a solid wall, solve for the resulting velocities taking the limit:


$$\lim_{m_b \rightarrow \infty} v'_a = \frac{m_a v_a + m_b v_b + e m_b (v_b - v_a)}{m_a + m_b}$$

$$= \lim_{m_b \rightarrow \infty} \left[\frac{m_a v_a}{m_a + m_b} + \frac{m_b v_b}{m_a + m_b} + \frac{e m_b (v_b - v_a)}{m_a + m_b} \right]$$

$$= [0 + v_b + e(v_b - v_a)]$$

Since mass b is immoveable, its velocity is zero, so that a 's post-collision velocity is:

$$v'_a = -e v_a$$



mjb - July 31, 2021

11


Collisions – Experimentally Determining the Coefficient of Restitution

12

Velocities are hard to measure live, but distances are not.

So, drop the object from a height h , and measure its bounce to a height h' :

Before the bounce:	After the bounce:
$v_2^2 = 0^2 + 2gh$	$0^2 = v'^2 - 2gh'$
$v = \sqrt{2gh}$	$v' = \sqrt{2gh'}$
$ v' = e v $	
<div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 0 auto;"> $e = \frac{v'}{v} = \frac{\sqrt{2gh'}}{\sqrt{2gh}} = \sqrt{\frac{h'}{h}}$ </div>	




mjb - July 31, 2021

12

Collisions – Some Coefficients of Restitution of Balls Bounced on a Concrete Surface

Ball Material	CoR
range golf ball	0.858
tennis ball	0.712
billiard ball	0.804
hand ball	0.752
wooden ball	0.603
steel ball bearing	0.597
glass marble	0.658
ball of rubber bands	0.828
hollow, hard plastic ball	0.688

<http://hypertextbook.com/facts/2006/restitution.shtml>



mjb – July 31, 2021

13


The Physics of Collisions – Totally Plastic Collisions

$$v'_a = \frac{m_a v_a + m_b v_b + e m_b (v_b - v_a)}{m_a + m_b}$$

$$v'_b = \frac{m_a v_a + m_b v_b - e m_a (v_b - v_a)}{m_a + m_b}$$

If $e=0$, then the two objects stick together and end up with the same resulting velocity:

$$v'_a = v'_b = \frac{m_a v_a + m_b v_b}{m_a + m_b}$$




mjb – July 31, 2021

14

One of my Jury Duties: Two vehicles collide. One is very damaged, the other hardly at all. What happened? Who's right?


Very damaged



B
25 mph

→

Hardly damaged



A
0 mph

$v_a = 0$, mph
 $v_b = 25$ mph = 11.2 m/sec
 $m_a = 2.0$
 $m_b = 1.0$
 $e = .30$


$$v'_a = \frac{v_b + e v_b}{m_a + m_b} = \frac{11.2(1.3)}{3} = 4.9 \text{ m/sec}$$

$$v'_b = \frac{m_b v_b - e m_a v_b}{m_a + m_b} = \frac{11.2(1 - .3 * 2)}{3} = 1.5 \text{ m/sec}$$

How much velocity change did each car undergo? (Δv is an approximate measure of damage.)

$\Delta v_a = 4.9 - 0 = 4.9 \text{ m/sec} = 11.0 \text{ mph}$

$\Delta v_b = 1.5 - 11.2 = -9.7 \text{ m/sec} = -21.7 \text{ mph}$



mjb – July 31, 2021

15

The Physics of Collisions – Totally Elastic Collisions


What happens when $e=1$? The two fundamental equations are

$$m_b v'_b + m_a v'_a = m_b v_b + m_a v_a$$

$$v'_b - v'_a = -e(v_b - v_a) = (v_a - v_b)$$

Rearranging:

$$m_a (v'_a - v_a) = m_b (v_b - v'_b)$$

$$v'_a + v_a = v_b + v'_b$$


mjb – July 31, 2021

16

The Physics of Collisions – Elastic Collisions

17

$$m_a (v'_a - v_a) = m_b (v_b - v'_b)$$

$$v'_a + v_a = v_b + v'_b$$

Then, multiplying the two together gives:

$$m_a v_a'^2 - m_a v_a^2 = m_b v_b^2 - m_b v_b'^2$$

Or:

$$\frac{1}{2} m_b v_b'^2 + \frac{1}{2} m_a v_a'^2 = \frac{1}{2} m_b v_b^2 + \frac{1}{2} m_a v_a^2$$

This shows that energy is conserved when the Coefficient of Restitution is 1.0




Oregon State
University
Computer Graphics

mjb - July 31, 2021

17

The Physics of Collisions – Oblique Impacts

18



Oblique Impacts are then handled by using vector math to determine the direct and tangential velocity components with respect to the **Line of Impact**.

The direct components are changed using the equations we just derived.

The tangential components are left unchanged. (This assumes no friction.)

The new components are then combined to produce the resulting velocity vectors.



Oregon State
University
Computer Graphics



mjb - July 31, 2021

18