1

Vectors

Oregon State University

Mike Bailey
mjb@cs.oregonstate.edu





1

2

Vectors have Direction and Magnitude



Magnitude: $\|V\| = \sqrt{V_x^2 + V_y^2 + V_z^2}$

2

3

A Vector Can Also Be Defined as the Positional Difference Between Two Points



$$(V_x, V_y, V_z) = (Q_x - P_x, Q_y - P_y, Q_z - P_z)$$

4

Unit Vectors have a Magnitude = 1.0



$$\|V\| = \sqrt{V_x^2 + V_y^2 + V_z^2}$$

$$\hat{V} = \frac{V}{\|V\|}$$

The circumflex (^) tells us this is a unit vector

5

Dot Product



$$A \bullet B = (A_x B_x + A_y B_y + A_z B_z) = \|A\|\|B\| \cos\theta$$

Because it produces a scalar result (i.e., a single number), this is also called the *Scalar Product*

6

A Physical Interpretation of the Dot Product



The amount of the force accelerating the car along the road is "how much of *F* is in the horizontal direction?"

$$F_{road} = F \cos\theta$$

This is easy to see in 2D, but a 3D version of the same problem is trickier.

A Physical Interpretation of the Dot Product

7



The amount of the force accelerating the car along the road is "how much of *F* is in the R direction?"

$$F_{road} = F\cos\theta = F \bullet \hat{R}$$

Oregon State University Computer Graphics

mjb – September 4, 2021

7

Even Trickier…

8



$$F_{road} = F\cos\theta = F \bullet \hat{R}$$

Oregon State University Computer Graphics

mjb – September 4, 2021

8

Generalizing How Much of A Lives in the B Direction

9



$$A \cdot B = \|A\|\|B\|cos\theta$$

$$A \cdot \hat{B} = \|A\|cos\theta$$

which is the length of the projection of A onto the B line

So, how much of A lives in the $\hat{B}$ direction is that magnitude times the B unit vector:

$$(A \cdot \hat{B})\ \hat{B}$$

Oregon State University Computer Graphics

mjb – September 4, 2021

9

Generalizing How Much of A Lives Perpendicular to the B Direction

10



From the previous slide, how much of A lives in the $\hat{B}$ direction is :

$$(A \cdot \hat{B})\ \hat{B}$$

That, plus the perpendicular vector equals A, so that how much of A is perpendicular to the $\hat{B}$ direction is:

$$A - (A \cdot \hat{B})\ \hat{B}$$

Oregon State University Computer Graphics

mjb – September 4, 2021

10

Dot Products are Commutative

11

$$A \bullet B = B \bullet A$$

Dot Products are Distributive

$$A \bullet (B + C) = (A \bullet B) + (A \bullet C)$$

Oregon State University Computer Graphics

mjb – September 4, 2021

11

The Perpendicular to a 2D Vector

12

If $V = (x, y)$

then $V_\perp = (-y, x)$

You can tell that this is true because

$$V \bullet V_\perp = (x, y) \bullet (-y, x) = -xy + xy = 0 = cos90°$$

Oregon State University Computer Graphics

mjb – September 4, 2021

12

Cross Product

13



$$A \times B = (A_y B_z - A_z B_y, A_z B_x - A_x B_z, A_x B_y - A_y B_x)$$

$$\|A \times B\| = \|A\| \|B\| \sin\theta$$

Because it produces a vector result (i.e., three numbers), this is also called the *Vector Product*

Oregon State University Computer Graphics

mjb – September 4, 2021

13

The Perpendicular Property of the Cross Product

14

The vector $A \times B$ is both perpendicular to A and perpendicular to B



The Right-Hand-Rule Property of the Cross Product

Curl the fingers of your right hand in the direction that starts at A and heads towards B. Your thumb points in the direction of AxB.

Oregon State University Computer Graphics

mjb – September 4, 2021

14

Cross Products are *Not* Commutative

15

$$A \times B = -B \times A$$



Cross Products are Distributive

$$A \times (B + C) = (A \times B) + (A \times C)$$

Oregon State University Computer Graphics

mjb – September 4, 2021

15

A CG Use for the Cross Product : Finding a Vector Perpendicular to a Plane (= the Surface Normal)

16



$$n = (R - Q) \times (S - Q)$$

Oregon State University Computer Graphics

mjb – September 4, 2021

16

A Use for the Cross Product : Finding a Vector Perpendicular to a Plane (= the Surface Normal) – This is used in CG Lighting

17



Oregon State University Computer Graphics

mjb – September 4, 2021

17

A Use for the Cross and Dot Products : Is a Point Inside a Triangle? – 3D (X-Y-Z) Version

18



Let:

$$n = (R - Q) \times (S - Q)$$

$$n_q = (R - Q) \times (P - Q)$$

$$n_r = (S - R) \times (P - R)$$

$$n_s = (Q - S) \times (P - S)$$

If $(n \bullet n_q), (n \bullet n_r), and (n \bullet n_s)$ are all positive, then P is inside the triangle QRS

Oregon State University Computer Graphics

mjb – September 4, 2021

18

Is a Point Inside a Triangle?
This can be simplified if you are in 2D (X-Y)

19



$E_{RS} = (P - R) \bullet (RS)_\perp$

where: $RS = (S_x - R_x, S_y - R_y)$

and: $(RS)_\perp = (R_y - S_y, S_x - R_x)$

Similarly,

$E_{SQ} = (P - S) \bullet (SQ)_\perp$

$E_{QR} = (P - Q) \bullet (QR)_\perp$

If E_{RS}, E_{SQ}, E_{QR} are all positive, then P is inside the triangle QRS

Oregon State University
Computer Graphics

mjb – September 4, 2021

19

A Use for the Cross Product : Finding the Area of a 3D Triangle

20



$$Area = \frac{1}{2} \cdot Base \cdot Height$$

$$Base = \|QR\|$$

$$Height = \|QS\| \sin\theta$$

$$Area = \frac{1}{2} \cdot \|QR\| \cdot \|QS\| \cdot \sin\theta = \frac{1}{2} \cdot \|(R - Q) \times (S - Q)\|$$

Oregon State University
Computer Graphics

mjb – September 4, 2021

20

Derivation of the Law of Sines

21



$$2 * Area(\Delta QRS) = \|(S - Q) \times (R - Q)\|$$

$$= rs \sin Q$$

But, the area is the same regardless of which two sides we use to compute it, so:

$$rs \sin Q = qs \sin R = qr \sin S$$

Dividing by *(qrs)* gives:

$$\frac{\sin Q}{q} = \frac{\sin R}{r} = \frac{\sin S}{s}$$

Oregon State University
Computer Graphics

mjb – September 4, 2021

21

Derivation of the Law of Cosines

22



$$s = R - Q$$

$$s^2 = \|R - Q\|^2$$

$$s^2 = (R - Q) \bullet (R - Q)$$

$$s^2 = [(R - S) + (S - Q)] \bullet [(R - S) + (S - Q)]$$

$$s^2 = [(R - S)(R - S)] + [(S - Q)(S - Q)] - 2(R - S) \bullet (S - Q)$$

$$s^2 = q^2 + r^2 - 2qr \cos S$$

Oregon State University
Computer Graphics

mjb – September 4, 2021

22

Distance from a Point to a Plane

23



In high school, you defined a plane by:
Ax + By + Cz + D = 0

It is more useful to define it by a point on the plane (Q) combined with the plane's normal vector (n_x, n_y, n_z)

If you want the familiar equation of the plane, it is:

$$\big((x, y, z) - (Q_x, Q_y, Q_z)\big) \bullet (n_x, n_y, n_z) = 0$$

which expands out to become the more familiar Ax + By + Cz + D = 0

The perpendicular distance from the point P to the plane is based on the plane equation:

$$d = (P - Q) \bullet \hat{n}$$

The dot product is answering the question "How much of (P-Q) is in the $\hat{n}$ direction?". Note that this gives a *signed distance*. If d > 0., then P is on the same side of the plane as the normal points. This is *very* useful.

Computer Graphics

mjb – September 4, 2021

23

Where does a line segment intersect an infinite plane?

24



The equation of the line segment is:

$$P = (1 - t)P_0 + tP_1$$

If point P is in the plane, then:

$$\big((P_x, P_y, P_z) - (Q_x, Q_y, Q_z)\big) \bullet (n_x, n_y, n_z) = 0$$

If we substitute the parametric expression for P into the plane equation, then the only thing we don't know in that equation is t. Solve it for t*. Knowing t* will let us compute the (x,y,z) of the actual intersection using the line equation. If t* has a zero in the denominator, then that tells us that t*=∞, and the line must be parallel to the plane.

This gives us the point of intersection with the infinite plane. We could now use the method covered a few slides ago to see if P lies inside a particular triangle.

Computer Graphics

mjb – September 4, 2021

24

Minimal Distance Between Two 3D Lines

25



The equation of the lines are : $P = P_0 + t \cdot v_p$ $\quad Q = Q_0 + t \cdot v_q$

The minimal distance vector between the two lines must be perpendicular to both

A vector between them that is perpendicular to both is: $v_\perp = v_p \times v_q$

We need to answer the question "How much of (Q_0-P_0) is in the $v_\perp$direction?".
To do this, we once again use the dot product:

$$d = (P_0 - Q_0) \bullet \hat{v}_\perp$$

Oregon State University
Computer Graphics

mjb – September 4, 2021

25

Another use for Dot Products : Force One Vector to be Perpendicular to Another Vector

26

Here, we want to force A to become perpendicular to B



The strategy is to get rid of the parallel component, leaving just the perpendicular

$$A = A_\parallel + A_\perp$$

So that: $A_\perp = A - A_\parallel$

But, $A_\parallel = (A \bullet \hat{B})\hat{B}$

So that $A' = A_\perp = A - (A \bullet \hat{B})\hat{B}$

Oregon State University
Computer Graphics

This is known as ***Gram-Schmidt orthogonalization***

mjb – September 4, 2021

26