




1


Resampling Scattered Data into a Regular Grid





Oregon State University
Mike Bailey
mjb@cs.oregonstate.edu






mjb - February 28, 2019

2

The Problem



National Doppler Radar Sites
Select radar location and click.

Alaska
Hawaii
South Korea
Puerto Rico


<http://www.ncdc.noaa.gov/nexradinv/>

Ofentimes data points are located irregularly, that is, they are not in nice, neat rectilinear grids.

This is called **Scattered Data**.

To use an Interpolated Color method, we need to triangulate the data so we can draw color-interpolated triangles.


To use Contours, we need to triangulate the data and re-sample it into a rectangular grid.



mjb - February 28, 2019


3

Once you have a Good Triangulation, You Can Superimpose any Data Grid You Want, and Re-sample the Data Values There



<http://www.ncdc.noaa.gov/nexradinv/>


We will see how to do that later in these notes.




mjb - February 28, 2019


4

Not all Triangulations are Created Equal: Which of These is Better, and Why?






<http://www.ncdc.noaa.gov/nexradinv/>



Long and Skinny Triangles




mjb - February 28, 2019

Three Steps

5

1. Fit a good set of triangles through the scattered points
2. Find out which triangle each new point is in
3. Interpolate within those triangles





mjb - February 28, 2019

1. Delauney Triangulation

6


This is an incremental algorithm, that is, you start with a "frame triangularization", and add a point at a time, adjusting the triangulation each time.


mjb - February 28, 2019

Adding a Point

7




1. Add a new point
2. Figure out which triangle it is in.
3. Create 3 new triangles by drawing lines from the new point to the 3 vertices of the bounding triangle




mjb - February 28, 2019

Look at each New Triangle, and Decide if it is Too Long and Skinny

8



2. Figure out which triangle it is in.
3. Create 3 new triangles by drawing lines from the new point to the 3 vertices of the bounding triangle
4. For each of the 3 new triangles, fit a circle through the 3 vertices (the new point, and the two existing points).



mjb - February 28, 2019

If it's Too Long and Skinny, Fix It 9

4. For each of the 3 new triangles, fit a circle through the 3 vertices (the new point, and the two existing points).
5. If the opposite point is inside the circle, then the circle is "too big", indicating that the triangle that was created is too long and skinny.
6. Delete the existing bounding edge, thus deleting two triangles.

Oregon State University
Computer Graphics

mjb - February 28, 2019

If it's Too Long and Skinny, Fix It 10

5. If the opposite point is inside the circle, then the circle is "too big", indicating that the triangle that was created is too long and skinny.
6. Delete the existing bounding edge, thus deleting two triangles.
7. Add a cross edge to make 2 new triangles

Oregon State University
Computer Graphics

mjb - February 28, 2019

A Very Slight Change in Point Location can affect the Triangulization 11

Oregon State University
Computer Graphics

mjb - February 28, 2019

A Use for the Cross and Dot Products :
Is a Point Inside a Triangle? - 3D (X-Y-Z) Version 12

Let:

$$n = (R - Q) \times (S - Q)$$

$$n_q = (R - Q) \times (P - Q)$$

$$n_r = (S - R) \times (P - R)$$

$$n_s = (Q - S) \times (P - S)$$

If $(n \bullet n_q), (n \bullet n_r),$ and $(n \bullet n_s)$
are all positive, then P is inside the triangle QRS


Oregon State University
Computer Graphics

Finding if a point is inside a triangle is used both in the Delaunay triangulization algorithm and in re-sampling to a new grid


mjb - February 28, 2019

3. Interpolating Data Values within a Triangle

13





Once we know the point is within a particular triangle, we need to interpolate within that triangle. Use a linear function:

$$S = Ax + By + C$$


mjb - February 28, 2019

**Think of the Scalar Function as Elevations
and Think of the Triangle Linear Interpolation Function as a Plane Being Fitted
on top of the Data Values**

14

mjb - February 28, 2019

**Since, at Vertices 0, 1, and 2, we know x, y, and s,
we can write 3 Equations with 3 Unknowns**


15

$$s_0 = Ax_0 + By_0 + C$$

$$s_1 = Ax_1 + By_1 + C$$

$$s_2 = Ax_2 + By_2 + C$$

or, in matrix form:

$$\begin{bmatrix} x_0 & y_0 & 1 \\ x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \end{bmatrix} \begin{Bmatrix} A \\ B \\ C \end{Bmatrix} = \begin{Bmatrix} s_0 \\ s_1 \\ s_2 \end{Bmatrix}$$



mjb - February 28, 2019

You can actually simplify it to 2 Equations with 2 Unknowns

16

$$\begin{aligned} s_0 &= Ax_0 + By_0 + C \\ s_1 &= Ax_1 + By_1 + C \\ s_2 &= Ax_2 + By_2 + C \end{aligned} \quad \rightarrow \quad \begin{aligned} s_1 - s_0 &= A(x_1 - x_0) + B(y_1 - y_0) \\ s_2 - s_0 &= A(x_2 - x_0) + B(y_2 - y_0) \end{aligned}$$

or, in matrix form:

$$\begin{bmatrix} x_1 - x_0 & y_1 - y_0 \\ x_2 - x_0 & y_2 - y_0 \end{bmatrix} \begin{Bmatrix} A \\ B \end{Bmatrix} = \begin{Bmatrix} s_1 - s_0 \\ s_2 - s_0 \end{Bmatrix}$$


mjb - February 28, 2019

17

Solve this 2x2 System in your Favorite Way – Cramer’s Rule Works Well


$$\begin{bmatrix} x_1 - x_0 & y_1 - y_0 \\ x_2 - x_0 & y_2 - y_0 \end{bmatrix} \begin{Bmatrix} A \\ B \end{Bmatrix} = \begin{Bmatrix} s_1 - s_0 \\ s_2 - s_0 \end{Bmatrix}$$

$$A = \frac{(s_1 - s_0)(y_2 - y_0) - (s_2 - s_0)(y_1 - y_0)}{(x_1 - x_0)(y_2 - y_0) - (x_2 - x_0)(y_1 - y_0)}$$

$$B = \frac{(x_1 - x_0)(s_2 - s_0) - (x_2 - x_0)(s_1 - s_0)}{(x_1 - x_0)(y_2 - y_0) - (x_2 - x_0)(y_1 - y_0)}$$

C is then computed by: $C = s_0 - Ax_0 - By_0$

Is it possible for A and/or B to compute to infinity?
What would that mean?




mjb – February 28, 2019

18

Interpolating Data Values within a Triangle: An Example

$S = Ax + By + C$




Vertex #0:
X = 0.
Y = 0.
S = 0.

Vertex #1:
X = 4.
Y = 0.
S = 12.

Vertex #2:
X = 3.
Y = 2.
S = 17.


(2,1)




mjb – February 28, 2019

19

The Delaunay Triangles can be used to Derive a Voronoi Diagram





Think of this as showing "Regions of Influence" around a Data Point



mjb – February 28, 2019

20


Voronoi Diagram:
Most of the Time, the Lines are the Perpendicular Bisectors of the Triangle Edges

mjb – February 28, 2019

Voronoi Regions of Influence

21




Oregon State
University
Computer Graphics

mjb - February 28, 2019