## Name: Answer Key

## Recitation time: \_\_\_\_\_

1. Let  $f : \mathbb{R}^3 \to \mathbb{R}^3$  be the linear map given by f(v) = Av where

$$A = \begin{bmatrix} 4 & -3 & -3 \\ 3 & -2 & -3 \\ -1 & 1 & 2 \end{bmatrix}.$$

Is f diagonalizable? If it is, express  $V = \mathbb{R}^3$  as a direct sum of one-dimensional invariant subspaces under f; then find a basis of V in which f is represented by a diagonal matrix. Solution: First find the characteristic polynomial of A:

$$det(A - \lambda I) = \begin{vmatrix} 4 - \lambda & -3 & -3 \\ 3 & -2 - \lambda & -3 \\ -1 & 1 & 2 - \lambda \end{vmatrix}$$
$$= (4 - \lambda) \begin{vmatrix} -2 - \lambda & -3 \\ 1 & 2 - \lambda \end{vmatrix} - 3 \begin{vmatrix} -3 & -3 \\ 1 & 2 - \lambda \end{vmatrix} + (-1) \begin{vmatrix} -3 & -3 \\ -2 - \lambda & -3 \end{vmatrix}$$
$$= (4 - \lambda)[(-2 - \lambda)(2 - \lambda) - (-3)(1)]$$
$$- 3[(-3)(2 - \lambda) - (-3)(1)]$$
$$- 3[(-3)(2 - \lambda) - (-3)(1)]$$
$$- [(-3)(-3) - (-3)(-2 - \lambda)]$$
$$= (4 - \lambda)(\lambda^2 - 1) - 3(3\lambda - 3) - (-3\lambda + 3)$$
$$= (4 - \lambda)(\lambda - 1)(\lambda + 1) - 3(3)(\lambda - 1) - (-3)(\lambda - 1))$$
$$= (\lambda - 1)[(4 - \lambda)(\lambda + 1) - 9 + 3]$$
$$= (\lambda - 1)[-\lambda^2 + 3\lambda - 2]$$
$$= (\lambda - 1)[-(\lambda - 1)(\lambda - 2)]$$
$$= -(\lambda - 1)^2(\lambda - 2)$$

Setting  $-(\lambda - 1)^2(\lambda - 2) = 0$  gives eigenvalues  $\lambda = 1$  and  $\lambda = 2$ .

We still do not know if f is diagonalizable. We now need to find the eigenvectors of A. This means solving the equation

$$(A - \lambda I)v = 0.$$

where  $v = (v_1, v_2, v_3)$ .

First let  $\lambda = -1$ . The augmented form of the above equation is

4 - 1	-3	-3	0		3	-3	-3	0
3	-2 - 1	-3	0	$\rightarrow$	3	-3	-3	0
1	1	2 - 1	0		-1	1	1	0

Row reduction gives

1	-1	$^{-1}$	0
0	0	0	0
0	0	0	0

Thus  $v_1 - v_2 - v_3 = 0$ . Adding  $v_2 + v_3$  to both sides gives  $v_1 = v_2 + v_3$ , so the eigenvectors v can be written as

$$v = \begin{bmatrix} v_2 + v_3 \\ v_2 \\ v_3 \end{bmatrix} = v_2 \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + v_3 \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$$

(continued on next page)

Therefore a basis for the eigenspace  ${\cal E}_1$  is

$$B_1 = \{(1, 1, 0), (1, 0, 1)\}$$

Now let  $\lambda = 2$ . We follow the same process of solving  $(A - \lambda I)v = 0$ :

$$\begin{bmatrix} 4-2 & -3 & -3 & 0 \\ 3 & -2-2 & -3 & 0 \\ -1 & 1 & 2-2 & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 2 & -3 & -3 & 0 \\ 3 & -4 & -3 & 0 \\ -1 & 1 & 0 & 0 \end{bmatrix}$$

Row reduction gives

$$\begin{bmatrix} 1 & 0 & 3 & 0 \\ 0 & 1 & 3 & 0 \\ -1 & 1 & 0 & 0 \end{bmatrix}$$

so we get the system of equations

$$v_1 + 3v_3 = 0$$
  
 $v_2 + 3v_3 = 0.$ 

This means that  $v_1 = v_2 = -3v_3$ , so the eigenvector v can be written as

$$v = \begin{bmatrix} -3v_3 \\ -3v_3 \\ v_3 \end{bmatrix} = v_3 \begin{bmatrix} -3 \\ -3 \\ 1 \end{bmatrix}.$$

A basis for  $E_2$  is

$$B_2 = \{(-3, -3, 1)\}$$

Since  $\dim(E_1) + \dim(E_2) = 2 + 1 = 3 = \dim(V)$  the linear map f is diagonalizable. Under the basis

$$B = B_1 \cup B_2 = \{(1, 1, 0), (1, 0, 1), (-3, -3, 1)\} = \{w_1, w_2, w_3\}$$

the map f is represented by the diagonal matrix

$$[f]_{B,B} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix},$$

and

$$V = \mathbb{R}^3 = \mathbb{R}w_1 \oplus \mathbb{R}w_2 \oplus \mathbb{R}w_3$$

where each  $\mathbb{R}w_i$  is an invariant subspace under f.

**2.** Let  $F : \mathbb{R}^{\infty} \to \mathbb{R}^{\infty}$  be the left-shift operator

$$F(x_1, x_2, x_3, \dots) = (x_2, x_3, x_4, \dots).$$

Find all eigenvalues and eigenvectors of F.

**Solution:** Here we are working with an infinite dimensional vector space, so we cannot simply find the eigenvectors of a matrix. We must use the definition of the eigenvector.

The eigenvectors of F are the vectors  $v = (x_1, x_2, ...) \in \mathbb{R}^\infty$  such that  $F(v) = \lambda v$  for some  $\lambda \in \mathbb{R}$ . In this case we have

$$(x_2, x_3, x_4, \dots) = \lambda(x_1, x_2, x_3, \dots)$$
$$= (\lambda x_1, \lambda x_2, \lambda x_3, \dots)$$

 $\mathbf{SO}$ 

$$x_2 = \lambda x_1$$
$$x_3 = \lambda x_2$$
$$x_4 = \lambda x_3$$
$$\vdots$$

We can now write every coordinate in terms of  $x_1$ :

$$\begin{aligned} x_2 &= \lambda x_1 \\ x_3 &= \lambda x_2 = \lambda (\lambda x_1) = \lambda^2 x_1 \\ x_4 &= \lambda x_3 = \lambda (\lambda^2 x_1) = \lambda^3 x_1 \\ \vdots \end{aligned}$$

so the eigenvector v looks like

$$v = (x_1, \lambda x_1, \lambda^2 x_1, \lambda^3 x_1, \dots)$$
$$= x_1(1, \lambda, \lambda^2, \lambda^3, \dots).$$

There is no restriction on  $\lambda$  in our work above, so every element  $\lambda \in \mathbb{R}$  is an eigenvalue. The eigenspace associated to the eigenvalue  $\lambda$  is

$$E_{\lambda} = \{x_1(1,\lambda,\lambda^2,\lambda^3,\dots) : x_1 \in \mathbb{R}\}\$$

which has basis

$$B_{\lambda} = \{(1, \lambda, \lambda^2, \lambda^3, \dots)\}.$$

**3.** Let  $g: \mathbb{C}^2 \to \mathbb{C}^2$  be the linear map given by g(v) = Bv where

$$B = \begin{bmatrix} i & 1\\ 2 & 1+i \end{bmatrix}.$$

(a) Find the eigenvalues of g. Why does this show that g is diagonalizable?Solution: Calculate the characteristic polynomial:

$$det(B - \lambda I) = \begin{vmatrix} i - \lambda & 1 \\ 2 & 1 + i - \lambda \end{vmatrix}$$
$$= (i - \lambda)(1 + i - \lambda) - 2$$
$$= \lambda^2 - (1 + 2i)\lambda + (-3 + i)$$

To solve  $det(A - \lambda I) = 0$  we can use the quadratic formula:

$$\lambda = \frac{(1+2i) \pm \sqrt{[-(1+2i)]^2 - 4(-3+i)}}{2}$$
$$= \frac{(1+2i) \pm \sqrt{-3 + 4i + 12 - 4i)}}{2}$$
$$= \frac{(1+2i) \pm \sqrt{9}}{2}$$
$$= \frac{(1+2i) \pm 3}{2}$$

 $\mathbf{SO}$ 

$$\lambda = 2 + i$$
 or  $\lambda = -1 + i$ .

Since  $\mathbb{C}^2$  is 2-dimensional and g has two distinct roots, this means that g is diagonalizable.

(b) Find a basis of  $\mathbb{C}^2$  in which g is represented by a diagonal matrix.

**Solution:** To find the eigenvectors, solve  $(A - \lambda I)v = 0$  where  $v = (v_1, v_2) \in \mathbb{C}^2$ . First let  $\lambda = 2 + i$ :

$$\begin{bmatrix} i - (2+i) & 1 & 0 \\ 2 & 1+i - (2+i) & 0 \end{bmatrix} \xrightarrow{\text{simplify}} \begin{bmatrix} -2 & 1 & 0 \\ 2 & -1 & 0 \end{bmatrix}$$
$$\xrightarrow{R_2 + R_1 \to R_2} \begin{bmatrix} -2 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

The first line of the reduced matrix gives  $-2v_1 + v_2 = 0$ , so  $v_2 = 2v_1$ . We can write the eigenvector v as

$$\begin{bmatrix} v_1\\2v_1 \end{bmatrix} = v_1 \begin{bmatrix} 1\\2 \end{bmatrix},$$

so a basis for  $E_{2+i}$  is

$$B_{2+i} = \{(1,2)\}$$

Now let  $\lambda = -1 + i$ :

$$\begin{bmatrix} i - (-1+i) & 1 & | & 0 \\ 2 & 1+i - (-1+i) & | & 0 \end{bmatrix} \xrightarrow{\text{simplify}} \begin{bmatrix} 1 & 1 & | & 0 \\ 2 & 2 & | & 0 \end{bmatrix}$$
$$\xrightarrow{R_2 - 2R_1 \to R_2} \begin{bmatrix} 1 & 1 & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix}$$

Then  $v_1 + v_2 = 0$ , so  $v_2 = -v_1$ . We can write the eigenvector v as

$$\begin{bmatrix} v_1 \\ -v_1 \end{bmatrix} = v_1 \begin{bmatrix} 1 \\ -1 \end{bmatrix},$$

so a basis for  $E_{-1+i}$  is

$$B_{-1+i} = \{(1, -1)\}$$

Let

$$B = \{(1,2), (1,-1)\} = \{w_1, w_2\}.$$

Then B is a basis for  $\mathbb{C}^2$  and the map g is represented by the diagonal matrix

$$[g]_{B,B} = \begin{bmatrix} 2+i & 0\\ 0 & -1+i \end{bmatrix},$$